

CHAPTER 5: TOPOLOGICAL FUNCTORS

## 1 Factorization structures

**Definition (5.1.1).** Let  $\mathbf{C}$  be a category.

- (i) If  $E$  is a class of  $\mathbf{C}$ -morphisms which is closed under composition with isomorphisms and  $M$  is a conglomerate of sources in  $\mathbf{C}$  which is closed under composition with isomorphisms, then the pair  $(E, M)$  is called a *factorization structure* on  $\mathbf{C}$  provided that the following are satisfied:
  - (a) For each source  $(X, (f_i)_{i \in I})$  there exists  $e: X \rightarrow Y$  in  $E$  and  $(Y, (m_i)_{i \in I})$  in  $M$  such that  $f_i = m_i \circ e$  for each  $i \in I$ ; briefly each source has  $(E, M)$ -factorization.
  - (b) For any two  $\mathbf{C}$ -morphisms  $f$  and  $e$  and any two sources  $(Y, (m_i)_{i \in I})$  and  $(Z, (f_i)_{i \in I})$  in  $\mathbf{C}$  such that  $e \in E$   $(Y, (m_i)_{i \in I}) \in M$  and  $f_i \circ e = m_i \circ f$  for each  $i \in I$ , there exists a unique  $\mathbf{C}$ -morphism  $g: Z \rightarrow Y$  such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{e} & Z \\ f \downarrow & \swarrow g & \downarrow f_i \\ Y & \xrightarrow{m_i} & X_i \end{array}$$

commutes for each  $i \in I$ ; briefly:  $\mathbf{C}$  satisfies the  $(E, M)$ -diagonalization property.

- (ii)  $\mathbf{C}$  is called  $(E, M)$ -category provided that  $(E, M)$  is a factorization structure on  $\mathbf{C}$ .

Let  $\mathbf{C}$  be a topological category or  $\mathbf{C} = \mathbf{Haus}$ .  $E$  consists of all extremal epimorphisms and  $M$  consists of all mono-sources in  $\mathbf{C}$ . Then  $(E, M)$  is a factorization structure on  $\mathbf{C}$ .

**Theorem (5.1.4).** Let  $\mathbf{C}$  be a category and  $(E, M)$  a factorization structure on  $\mathbf{C}$ . Then the following are satisfied:

- (i)  $(E, M)$ -factorizations are uniquely determined (up to isomorphisms).
- (ii)  $E \cap M$  is the class of all  $\mathbf{C}$ -isomorphisms.
- (iii) Every extremal source in  $\mathbf{C}$  belongs to  $M$ .
- (iv) If  $f, g$  and  $h$  are  $\mathbf{C}$ -morphisms such that  $h = g \circ f$ , then the following are satisfied:
  - (a) If  $h \in E$  and  $f$  is a  $\mathbf{C}$ -epimorphism, then  $g \in E$ .
  - (b)  $f \in E$  and  $g \in E$  imply  $h \in E$ , i.e.  $E$  is closed under composition.
- (v) If  $(X, (f_i)_{i \in I})$  is a source in  $\mathbf{C}$  and  $(X, (g_j)_{j \in J}), (Z_j, (k_{j_i})_{i \in I_j})_{j \in J}$  is a factorization of  $(X, (f_i)_{i \in I})$ , then the following hold:
  - (a)  $(X, (f_i)_{i \in I}) \in M$  implies  $(X, (g_j)_{j \in J}) \in M$ .
  - (b)  $(X, (g_j)_{j \in J}) \in M$  and  $(Z_j, (k_{j_i})_{i \in I_j}) \in M$  for each  $j \in J$  imply  $(X, (f_i)_{i \in I}) \in M$ .

- (vi) If  $(X, (f_i)_{i \in I})$  is a source in  $\mathbf{C}$  and there is some  $J \subset I$  such that  $(X, (f_j)_{j \in J}) \in M$ , then  $(X, (f_i)_{i \in I}) \in M$ .
- (vii)  $E$  and  $M$  determine each other by the diagonalization property.

## 2 Definition and properties of topological functors

**Definition (5.2.1).** Let  $\mathbf{C}$  be a category supplied with a factorization structure  $(E, M)$ , let  $\mathbf{A}$  be any category and let  $T: \mathbf{A} \rightarrow \mathbf{C}$  be a functor.

- (i) A source  $(A, (f_i : A \rightarrow A_i)_{i \in I})$  in  $\mathbf{A}$  is called *T-initial* provided that for each source  $(B, (g_i : B \rightarrow A_i)_{i \in I})$  in  $\mathbf{A}$  and each  $\mathbf{C}$ -morphism  $f: T(A) \rightarrow T(B)$  such that  $T(f_i) \circ f = T(g_i)$  for each  $i \in I$ , there exists a unique  $\mathbf{A}$ -morphism  $\bar{f}: B \rightarrow A$  with  $T(\bar{f}) = f$  and  $f_i \circ \bar{f} = g_i$  for each  $i \in I$ .
- (ii) A source  $(A, (f_i : A \rightarrow A_i)_{i \in I})$  in  $\mathbf{A}$  *T-lifts* a source  $(X, (g_i : X \rightarrow T(A_i))_{i \in I})$  in  $\mathbf{C}$  provided that there exists an isomorphism  $h: X \rightarrow T(A)$  in  $\mathbf{C}$  with  $T(f_i) \circ h = g_i$  for each  $i \in I$ .
- (iii)  $T$  is called  *$(E, M)$ -topological* provided that for each family  $(A_i)_{i \in I}$  of  $\mathbf{A}$ -objects and each source  $(X, (m_i : X \rightarrow T(A_i))_{i \in I})$  in  $M$ , there exists a *T-initial* source  $(A, (f_i : A \rightarrow A_i)_{i \in I})$  in  $\mathbf{A}$  which *T-lifts*  $(X, (m_i)_{i \in I})$ .
- (iv)  $T$  is called *absolutely topological* provided that  $T$  is topological for any factorization structure  $(E, M)$  on  $\mathbf{C}$ .

**Theorem (5.2.4).** Let  $\mathbf{C}$  be a category supplied with a factorization structure  $(E, M)$  and let  $T: \mathbf{A} \rightarrow \mathbf{C}$  be an  $(E, M)$ -topological functor. Then  $T$  is faithful.

**Theorem (5.2.5).** Let  $\mathbf{C}$  be an  $(E, M)$ -category and  $T: \mathbf{A} \rightarrow \mathbf{C}$  an  $(E, M)$ -topological functor. If  $E_T$  denotes the class of all morphisms  $f$  in  $\mathbf{A}$  with  $T(f) \in E$  and  $M_T$  denotes the conglomerate of all *T-initial* sources  $(A, (f_i)_{i \in I})$  in  $\mathbf{A}$  with  $(T(A), (T(f_i))_{i \in I}) \in M$ , then  $\mathbf{A}$  is an  $(E_T, M_T)$ -category.

**Theorem (5.2.8).** Let  $T: \mathbf{A} \rightarrow \mathbf{C}$  be an  $(E, M)$ -topological functor, let  $D: \mathbf{I} \rightarrow \mathbf{A}$  be a diagram and let  $(L, (l_i : L \rightarrow D(i))_{i \in |\mathbf{I}|})$  be a source in  $\mathbf{A}$ . Then the following are equivalent:

- (i)  $(L, (l_i)_{i \in |\mathbf{I}|})$  is a limit of  $D$ .
- (ii)  $(L, (l_i)_{i \in |\mathbf{I}|})$  is *T-initial* and  $(T(L), (T(l_i))_{i \in |\mathbf{I}|})$  is a limit of  $T \circ D$ .

**Corollary (5.2.9).** If  $T: \mathbf{A} \rightarrow \mathbf{C}$  is an  $(E, M)$ -topological functor and  $\mathbf{C}$  is complete, then  $\mathbf{A}$  is complete.

**Proposition (5.2.10).** Let  $T: \mathbf{A} \rightarrow \mathbf{C}$  be an  $(E, M)$ -topological functor. Then for each family  $(A_i)_{i \in I}$  of  $\mathbf{A}$ -objects and each sink  $((f_i : T(A_i) \rightarrow X)_{i \in I}, X)$  in  $\mathbf{C}$ , there exists a sink  $((t_i : A_i \rightarrow A)_{i \in I}, A)$  in  $\mathbf{A}$  and a morphism  $e: X \rightarrow T(A)$  in  $E$  such that  $T(t_i) = e \circ f_i$  for each  $i \in I$  and such that the following condition is satisfied:

(F) For each sink  $((g_i : A_i \rightarrow B)_{i \in I}, B)$  in  $\mathbf{A}$  and each morphism  $g: X \rightarrow T(B)$  with  $T(g_i) = g \circ f_i$  for each  $i \in I$ , there exists a morphism  $k: A \rightarrow B$  with  $T(k) \circ e = g$ .

**Theorem (5.2.11).** *Let  $T: \mathbf{A} \rightarrow \mathbf{C}$  be an  $(E, M)$ -topological functor and let  $D: \mathbf{I} \rightarrow \mathbf{A}$  be a diagram such that  $T \circ D$  has a colimit. Then  $D$  has a colimit.*

**Corollary (5.2.12).** *Let  $T: \mathbf{A} \rightarrow \mathbf{C}$  be an  $(E, M)$ -topological functor. If  $\mathbf{C}$  is cocomplete, then  $\mathbf{A}$  is cocomplete.*

### 3 Initially structured categories

A functor  $F: \mathbf{A} \rightarrow \mathbf{B}$  is called *amnestic* provided that any  $\mathbf{A}$ -isomorphism  $f$  is an  $\mathbf{A}$ -identity iff  $F(f)$  is a  $\mathbf{B}$ -identity.

A functor  $F: \mathbf{A} \rightarrow \mathbf{B}$  is called *transportable* provided that for each  $\mathbf{A}$ -object  $A$ , each  $\mathbf{B}$ -object  $B$  and each isomorphism  $q: B \rightarrow F(A)$ , there exists a unique  $\mathbf{A}$ -object  $C$  and isomorphism  $\bar{q}: C \rightarrow A$  with  $F(\bar{q}) = q$ .

An object  $X$  in a category  $\mathbf{C}$  is called *terminal* provided that for every object  $Y$  in  $\mathbf{C}$ , the set of morphisms  $[Y, X]_{\mathbf{C}}$  is a singleton.

**Definition (5.3.1).** A pair  $(\mathbf{A}, T)$  is called an *initially structured* category provided that  $\mathbf{A}$  is a category and  $T: \mathbf{A} \rightarrow \mathbf{Set}$  is a functor which is amnestic and transportable such that the following hold:

- (i)  $T$  is (epi, mono-source)-topological.
- (ii)  $T$  has small fibres, i.e. for each  $X \in \mathbf{Set}$   $\{A \in |\mathbf{A}| : T(A) = X\}$  is a set.
- (iii) There is precisely one object  $P$  in  $\mathbf{A}$  (up to isomorphism) such that  $T(P)$  is a terminal separator in  $\mathbf{Set}$ , i.e.  $T(P)$  is a singleton.

**Remark (5.3.3).** Obviously the condition (i) can be replaced by:

Every source  $(X, (f_i: X \rightarrow T(A_i))_{i \in I})$  in  $\mathbf{Set}$  has an (epi, mono-source) factorization

$$\begin{array}{ccc} X & \xrightarrow{f_i} & T(A_i) \\ & \searrow e & \nearrow T(g_i) \\ & T(B) & \end{array}$$

**Theorem (5.3.4).** *Let  $(\mathbf{A}, T)$  be an initially structured category. Then the following are satisfied:*

- (i)  $T$  is faithful.
- (ii) A source  $(A, (f_i: A \rightarrow D_i)_{i \in I})$  in  $\mathbf{A}$  is a limit of a diagram  $D: \mathbf{I} \rightarrow \mathbf{A}$  with  $|\mathbf{I}| = I$  if and only if this source is  $T$ -initial and  $(T(A), (T(f_i): T(A) \rightarrow T(D_i))_{i \in I})$  is limit of  $T \circ D$ .
- (iii) For any sink  $((f_i: D_i \rightarrow A)_{i \in I, A})$  in  $\mathbf{A}$  and an epimorphism  $e: X \rightarrow T(A)$  with  $e \circ f_i = T(a_i)$  for each  $i \in I$  such that the following condition (F) is satisfied: For each sink  $((b_i: D_i \rightarrow B)_{i \in I}, B)$  in  $\mathbf{A}$  and each morphism  $d: X \rightarrow T(B)$  there exists a (unique) morphism  $c: A \rightarrow B$  such that  $T(c) \circ e = d$ .

(iv)  $\mathbf{A}$  is complete and cocomplete.

**Proposition (5.3.5).** Let  $(\mathbf{A}, T)$  be an initially structured category. Any sink  $((f_i: A_i \rightarrow C)_{i \in I}, C)$  in  $\mathbf{A}$  has

(i) a factorization

$$\begin{array}{ccc} A_i & \xrightarrow{f_i} & C \\ a_i \searrow & \nearrow c & \\ & A & \end{array}$$

such that  $T(c)$  is an isomorphism and  $((a_i: A_i \rightarrow A)_{i \in I}, A)$  is  $T$ -final (i.e. for any sink  $((b_i: A_i \rightarrow B)_{i \in I}, B)$  and any morphism  $f: T(A) \rightarrow T(B)$  with  $f \circ T(a_i) = T(b_i)$  for each  $i \in I$ , there exists a unique morphism  $\bar{f}: A \rightarrow B$  with  $T(\bar{f}) = f$  and  $\bar{f} \circ a_i = b_i$  for each  $i \in I$ ), and

(ii) a factorization  $f_i = c \circ a_i$  where  $c$  is a monomorphism and  $((a_i: A_i \rightarrow A)_{i \in I}, A)$  is a  $T$ -final epi-sink.

**Proposition (5.3.6).** Let  $(\mathbf{A}, T)$  be an initially structured category. Then the following are satisfied:

- (i) The object  $P$  in  $\mathbf{A}$  given by  $IS_3$  (definition of initially structured category, (iii)) is terminal and a separator in  $\mathbf{A}$ .
- (ii) If  $X$  and  $Y$  are  $\mathbf{A}$ -objects and  $g: T(X) \rightarrow T(Y)$  is a constant morphism, then there exists a unique  $\mathbf{A}$ -morphism  $f: X \rightarrow Y$  with  $T(f) = g$ .

**Definition (5.3.7).** Let  $(\mathbf{A}, T)$  be an initially structured category. An  $\mathbf{A}$ -morphism  $f: A \rightarrow B$  is called a

- (i)  $T$ -embedding provided that  $(A, f: A \rightarrow B)$  is a  $T$ -initial mono-source.
- (ii)  $T$ -quotient map provided that  $(A, f: A \rightarrow B)$  is a  $T$ -final epi-sink.

**Proposition (5.3.8).** Let  $(\mathbf{A}, T)$  be an initially structured category. Then the following are satisfied:

- (i)  $f$  is an  $\mathbf{A}$ -monomorphism if and only if  $T(f)$  is a monomorphism in  $\mathbf{Set}$ .
- (ii)  $f$  is an  $\mathbf{A}$ -epimorphism whenever  $T(f)$  is an epimorphism in  $\mathbf{Set}$ .
- (iii) Every extremal monomorphism in  $\mathbf{A}$  is a  $T$ -embedding.

**Proposition (5.3.10).** Let  $(\mathbf{A}, T)$  be an initially structured category. Then the following are satisfied:

- (i) If  $((f_i: A_i \rightarrow A)_{i \in I}, A)$  is a  $T$ -final epi-sink in  $\mathbf{A}$ , then  $((T(f_i): T(A_i) \rightarrow T(A))_{i \in I}, T(A))$  is an epi-sink in  $\mathbf{Set}$ .
- (ii) A sink in  $\mathbf{A}$  is an extremal epi-sink if and only if it is a  $T$ -final epi-sink.

**Proposition (5.3.11).** *For every  $\mathbf{A}$ -morphism  $f: A \rightarrow B$  in an initially structured category  $(\mathbf{A}, T)$  the following are equivalent:*

- (i)  $f$  is a  $T$ -quotient map.
- (ii)  $f$  is a regular epimorphism.
- (iii)  $f$  is an extremal epimorphism.

**Proposition (5.3.13).** *Every initially structured category  $(\mathbf{A}, T)$  is well-powered.*

**Proposition (5.3.15).** *Let  $(\mathbf{A}, T)$  be an initially structured category. Then  $\mathbf{A}$  is a category with a factorization structure  $(E, M)$  where  $E$  consists of all extremal epimorphisms and  $M$  of all mono-sources.*

**Theorem (5.3.16).** *Every  $E$ -reflective (full and isomorphism-closed) subcategory of an initially structured category  $(\mathbf{A}, T)$  is initially structured provided that  $E$  consists of all  $\mathbf{A}$ -morphisms  $f$  for which  $T(f)$  is an epimorphism in  $\mathbf{Set}$ .*

**Remark (5.3.17).** Every extremal epireflective (resp. epireflective) full and isomorphism-closed subcategory of an initially structure (resp. topological) category is initially structured.

**Proposition (5.3.18).** *For each  $T$ -final epi-sink  $(f_i: A_i \rightarrow A)_{i \in I}$  in an initially structured category, there is a set  $J \subset I$  such that  $(f_j: A_j \rightarrow A)_{j \in J}$  is likewise a  $T$ -final epi-sink.*

**Proposition (5.3.19).** *Let  $(\mathbf{A}, T)$  and  $(\mathbf{A}', T')$  be initially structured categories and let  $F: \mathbf{A} \rightarrow \mathbf{A}'$  be a functor preserving colimits. If  $(f_i: A_i \rightarrow A)$  is a  $T$ -final epi-sink in  $\mathbf{A}$ , then  $(T(f_i): T(A_i) \rightarrow T(A))$  is a  $T'$ -final epi-sink in  $\mathbf{A}'$ .*

**Theorem (5.3.20).** *Let  $(\mathbf{A}, T)$  be an initially structured category.  $\mathbf{A}$  is cartesian closed if and only if  $A \times -$  preserves  $T$ -final epi-sinks for each  $A \in |\mathbf{A}|$ .*

**Theorem (5.3.22).** *Every extremal epireflective (full and isomorphism-closed) subcategory  $\mathbf{B}$  of an initially structured cartesian closed category  $(\mathbf{A}, T)$  is cartesian closed.*

**Theorem (5.3.24).** *For a concrete category  $\mathbf{C}$  (in the sense of 1.1.1) the following are equivalent:*

- (i)  $\mathbf{C}$  is initially structured.
- (ii)  $\mathbf{C}$  is an epireflective (full and isomorphism-closed) subcategory of a topological category.
- (iii)  $\mathbf{C}$  is an extremal epireflective (full and isomorphism-closed) subcategory of a topological category.